

Superheavy Dark Matter with Discrete Gauge Symmetries

K. Hamaguchi^{a)}, Yasunori Nomura^{a)†}, and T. Yanagida^{a,b)}

^{a)} *Department of Physics, University of Tokyo, Tokyo 113-0033, Japan*

^{b)} *RESCEU, University of Tokyo, Tokyo 113-0033, Japan*

Abstract

We show that there are discrete gauge symmetries protect naturally heavy X particles from decaying into the ordinary light particles in the supersymmetric standard model. This makes the proposal very attractive that the superheavy X particles constitute a part of the dark matter in the present universe. It is more interesting that there are a class of discrete gauge symmetries which naturally accommodate a long-lived unstable X particle. We find that in some discrete $\mathbf{Z}_{10}$ models, for example, a superheavy X particle has lifetime $\tau_X \simeq 10^{11} - 10^{26}$ years for its mass $M_X \simeq 10^{13} - 10^{14}$ GeV. This long lifetime is guaranteed by the absence of lower dimensional operators (of light particles) couple to the X . We briefly discuss a possible explanation for the recently observed ultra-high-energy cosmic ray events by the decay of this unstable X particle.

[†]Research Fellow of the Japan Society for the Promotion of Science.

The existence of invisible dark matter in the present universe is one of the central issues in particle physics as well as in astrophysics. Many particle candidates have been, in fact, proposed as constituents of the dark matter. It is usually thought that the dark-matter particles cannot be too heavy, since stable particles much heavier than ~ 1 TeV would easily overclose the universe if they have been once in thermal equilibrium [1]. Therefore, if one would consider superheavy dark matter, one must invoke, for example, a late-time entropy production to dilute sufficiently the abundance of such superheavy relic particles [2]. This may render this scenario unlikely.

Recently, the above problem has been naturally solved by the authors of [3, 4]. They have suggested that inflation in the early universe may generate a desirable amount of such a superheavy particle during the reheating epoch just after the end of inflation. Since the production mechanism for the superheavy particle involves only gravitational interactions, it is quite independent of detailed nature of the particle as well as processes of the reheating. This new observation seems very generic and thus it leads us to consider, in this paper, the proposal that most of the dark matter in the universe indeed consists of some superheavy particles [2]. The numerical calculation in ref. [3] shows that the desired abundance of the superheavy particle X is obtained when its mass M_X lies in the region $0.04 \lesssim M_X/H \lesssim 2$ with the Hubble parameter at the end of inflation $H \sim 10^{13}$ GeV. In this paper, we assume $M_X \simeq 10^{13} - 10^{14}$ GeV.

For the X particle to be a dominant component of the dark matter in the present universe, it must live longer than the age of the universe. Such stability of the superheavy X particle may be explained by imposing some symmetries. We consider here gauge (local) symmetries, since any global symmetries are broken explicitly by topological effects of gravity [5] and the X particle may no longer survive until the present.¹

There are two classes of gauge symmetries, continuous and discrete. We concentrate our discussion on discrete gauge symmetries [7], since analyses for continuous

¹ The lifetime of the X particle can be longer than the age of the universe if topological effects of gravity are extremely small [6].

(non-Abelian) gauge symmetries have been addressed in [8, 9]. A discrete gauge $\mathbf{Z}_2$ symmetry [10] (called R-parity) is already used to avoid the rapid proton decay due to dimension 4 operators in the supersymmetric (SUSY) standard model. We make our analysis in the framework of SUSY standard model introducing one superheavy X particle. We impose a $\mathbf{Z}_N$ gauge symmetry in addition to the R-parity and discuss if there are consistent discrete gauge symmetries that guarantee the long lifetime of the X particle required for the dark matter. We find various examples for such discrete symmetries. It may be very interesting that some of them naturally accommodate a superheavy X particle of mass $M_X \simeq 10^{13} - 10^{14}$ GeV with long lifetime $\tau_X \simeq 10^{11} - 10^{26}$ years. We briefly comment on a possible explanation for the recently observed ultra-high-energy cosmic ray events by the decay of this unstable superheavy particle.

First of all, N should be even so that the mass term for X is allowed in superpotential as

$$W = \frac{M_X}{2} X^2. \quad (1)$$

Here, we assume that X carries $N/2$ of the $\mathbf{Z}_N$ charge. Its R-parity could be even or odd. $\mathbf{Z}_N$ charges for the SUSY standard-model particles are given by the following discussion.

- (i) All terms in superpotential present in the SUSY standard model must be allowed by $\mathbf{Z}_N$.
- (ii) All anomalies for $\mathbf{Z}_N$ should vanish (cancellation with the Green-Schwarz term [11] will be given later).

As for R-parity for the light particles we adopt the charges given in ref. [10].

The first condition (i) reduces the number of independent $\mathbf{Z}_N$ charges for the light particles as shown in Table 1. Here, we have chosen the $\mathbf{Z}_N$ charge for a Higgs doublet H_u to be zero. This can be done by using a gauge rotation of $U(1)_Y$ in the standard model without a loss of generality.

	Q	$\bar{u}$	d	l	$\bar{e}$	H_u	H_d	X
$\mathbf{Z}_N$	m	$-m$	$-m$	p	$-p$	0	0	$N/2$
R	—	—	—	—	—	+	+	+ or —

Table 1: $\mathbf{Z}_N$ discrete charges for the SUSY standard-model particles and X . Here, $p, m = 1, \dots, N-1$. R denotes the R-parity. $Q, \bar{u}, \bar{d}, l$ and $\bar{e}$ denote $\text{SU}(2)_L$ -doublet quark, up-type antiquark, down-type antiquark, $\text{SU}(2)_L$ -doublet lepton and antilepton chiral multiplets. H_u and H_d are chiral multiplets for Higgs doublets.

From the second requirement (ii) we obtain a set of constraints [12]:

$$\begin{cases} 0 = \frac{1}{2}r'N \\ \frac{9}{2}m + \frac{3}{2}p = \frac{1}{2}r''N \\ 3p = \frac{1}{2}r'''N. \end{cases} \quad (r', r'', r''' \in \mathbf{Z}) \quad (2)$$

Here, p, m are $\mathbf{Z}_N$ charges for the SUSY standard-model particles given in Table 1. The first equation comes from the cancellation of $\{\mathbf{Z}_N\}\{\text{SU}(3)_C\}^2$ anomalies, the second from the cancellation of $\{\mathbf{Z}_N\}\{\text{SU}(2)_L\}^2$ anomalies, and the last from the cancellation of $\mathbf{Z}_N$ -gravitational anomalies. Notice that the anomaly-free condition for discrete gauge symmetries is much weaker than that for continuous ones [12].

These equations (2) are satisfied only if

$$\begin{cases} p = r_1 \left(\frac{N}{6} \right) \\ m = (2r_2 - r_1) \left(\frac{N}{18} \right). \end{cases} \quad (r_1, r_2 \in \mathbf{Z}) \quad (3)$$

We discard the trivial case, $p = m = 0$. Then, we find nontrivial discrete gauge symmetries,

$$\mathbf{Z}_2 : (p, m) = (1, 1), \quad (4)$$

$$\mathbf{Z}_6 : (p, m) = (1, 1), (2, 0), \quad (5)$$

$$\mathbf{Z}_{18} : (p, m) = (0, 4), (3, 1), (3, 5), (6, 2), (6, 4), (9, 1). \quad (6)$$

There are other solutions which satisfy eq. (3), but they are trivial embeddings of the above $\mathbf{Z}_2$, $\mathbf{Z}_6$ or $\mathbf{Z}_{18}$ in higher N , or charge conjugations or $\text{U}(1)_Y$ -gauge

equivalents of some of eqs. (4), (5) and (6). Notice that if the $\mathbf{Z}_N$ charge m is even the p is even, and if the m is odd the p is odd, since N is even (see the second eq. of (2)). Furthermore, the symmetry (4) is nothing but the matter parity, which is equivalent to the R-parity in the context of the SUSY standard model. Thus, nontrivial discrete gauge symmetries besides the R-parity are only the $\mathbf{Z}_6$ symmetry with charges in eq. (5) and $\mathbf{Z}_{18}$ symmetry with charges in eq. (6).²

Let us now discuss operators $\mathcal{O}$ and $\mathcal{O}'$, which consist only of the SUSY standard-model particles, defined in either super- or Kähler potentials as

$$W = \mathcal{O}X \quad , \quad K = \mathcal{O}'X. \quad (7)$$

The operators $\mathcal{O}$ and $\mathcal{O}'$ must carry $N/2 \pmod{N}$ of the $\mathbf{Z}_N$ charge to make super- and Kähler potentials W and K invariant under the $\mathbf{Z}_N$. If the $\mathbf{Z}_N$ charge m is even (in this case, the p is always even), the operators $\mathcal{O}$ and $\mathcal{O}'$ have always even $\mathbf{Z}_N$ charge, since all the light particles have even charges in this case. Thus, any operators $\mathcal{O}$ and $\mathcal{O}'$ consist of the light particles cannot have charge 3 or 9 ($= N/2$), and hence the X particle is completely stable for the even m .

If the m is odd (in this case, the p is always odd), we easily see that the total number of matter multiplets ($Q, \bar{u}, \bar{d}, l$ and $\bar{e}$ in Table 1) in the operators $\mathcal{O}$ and $\mathcal{O}'$ must be odd so that the $\mathcal{O}$ and $\mathcal{O}'$ carry 3 (9) of the $\mathbf{Z}_6$ ($\mathbf{Z}_{18}$) charge. On the other hand, we find from the $\mathbf{Z}_N$ charge assignment in Table 1 that any operators which have odd total number of matter multiplets carry always odd R-parity. This proves that the X particle with even R-parity is completely stable and can survive until the present as a superheavy dark matter. Notice that the condensation, $\langle H_u \rangle \neq 0$ and $\langle H_d \rangle \neq 0$, does not break any $\mathbf{Z}_N$. It is very surprising that anomaly-free condition (2) plays an essential role on deriving the conclusion that the discrete gauge symmetries $\mathbf{Z}_6$ and $\mathbf{Z}_{18}$ in eqs. (5) and (6) can naturally accommodate a stable X particle as long as X has even R-parity.³

² If one takes a basis where the $\mathbf{Z}_N$ charge for Q vanishes, both of the $\mathbf{Z}_6$ and $\mathbf{Z}_{18}$ become $\mathbf{Z}_6$. Therefore, precisely speaking, the anomaly-free discrete symmetry independent of the $U(1)_Y$ in the standard model is only $\mathbf{Z}_6$ besides the R-parity.

³ For the trivial case, $p = m = 0$, the X particle is obviously stable for any N , since all light particles do not have non-vanishing $\mathbf{Z}_N$ charges.

N	(p, m)	the lowest dimensional operator
6	(1, 1)	$\bar{u}dd \quad (\in \mathcal{O})$
18	(3, 5)	$\bar{d}\bar{d}\bar{d}ll \quad (\in \mathcal{O})$
18	(3, 1)	$\bar{u}\bar{u}\bar{u}\bar{e}\bar{e} \quad (\in \mathcal{O})$
18	(9, 1)	$lH_u \quad (\in \mathcal{O})$

Table 2: The lowest dimensional operators which X can couple to in the case that X has odd R-parity. If X has even R-parity, it is completely stable as explained in the text.

If the X has odd R-parity and the m is odd, it can couple to the light particles. The lowest dimensional operators in the $\mathbf{Z}_6$ and $\mathbf{Z}_{18}$ models are given in Table 2.⁴ We find that the lifetime of the X particle is shorter than 10^{-7} sec. The X particle with such a short lifetime, however, is astrophysically less interesting.

Let us turn to the case where the anomalies for $\mathbf{Z}_N$ are cancelled by the Green-Schwarz term [11]. As shown in ref. [12], we have the following constraints:

$$\begin{cases} 0 = \frac{1}{2}r'N + k\delta_{GS} \\ \frac{9}{2}m + \frac{3}{2}p = \frac{1}{2}r''N + k\delta_{GS} \\ 3p = \frac{1}{2}r'''N + 24\delta_{GS}, \end{cases} \quad (r', r'', r''' \in \mathbf{Z}) \quad (8)$$

where k is the Kac-Moody level of the $SU(3)_C$ and $SU(2)_L$ gauge groups, and δ_{GS} is a constant [12]. Here, we have set the Kac-Moody levels of $SU(3)_C$ and $SU(2)_L$ to be the same in view of the gauge coupling unification.

The solution of eq. (8) are

$$\begin{cases} p = \left(r_1 + \frac{24}{k}r_3\right) \left(\frac{N}{6}\right) \\ m = \left(2r_2 - r_1 - \frac{24}{k}r_3\right) \left(\frac{N}{18}\right), \end{cases} \quad (r_1, r_2, r_3 \in \mathbf{Z}) \quad (9)$$

and we have many independent discrete gauge symmetries which satisfy eq. (9). We have searched them up to $N = 18$, and in Table 3 we show all solutions, p and m , except for those in eqs. (4), (5) and (6). From the first and second eqs. of (8),

⁴ The dimension of the operator $\mathcal{O}$ consists of n chiral multiplets is the same as that of the operator $\mathcal{O}'$ consists of $n - 1$ chiral multiplets.

N	(p, m)
4	(1, 1)
8	(1, 5), (3, 7)
10	(1, 3), (2, 6), (3, 9), (4, 2)
12	(1, 1), (5, 1)
14	(1, 9), (3, 13), (5, 3)
16	(1, 5), (3, 15), (5, 9), (7, 3)

Table 3: Sets of solutions, p and m , for $N \leq 18$ which satisfy the anomaly-free constraints (8) relaxed in the presence of the Green-Schwarz term. For $N = 6$ and 18, see eqs. (5) and (6) in the text.

we find that if the $\mathbf{Z}_N$ charge m is odd the p is odd, and if the m is even the p is even, since N is even.

When $N/2$ is even, the $\mathbf{Z}_N$ charge m (and p) must be odd, otherwise it becomes a trivial extension of a lower N case. If the m (and p) is odd, the total number of matter multiplets in the operators $\mathcal{O}$ and $\mathcal{O}'$ must be even so that the $\mathcal{O}$ and $\mathcal{O}'$ have even $\mathbf{Z}_N$ charges. But, we see from the charge assignment in Table 1 that any operators which have even total number of matter multiplets carry always even R-parity. Thus, the X particle with odd R-parity is stable when $N/2$ is even.

On the contrary, when $N/2$ is odd, the m (and p) can be either odd or even. If the m (and p) is even, the $\mathcal{O}$ and $\mathcal{O}'$ never have odd ($= N/2$) $\mathbf{Z}_N$ charges, since all light particles have even $\mathbf{Z}_N$ charges in this case. Thus, the X particle can not couple to any operators $\mathcal{O}$ and $\mathcal{O}'$ consist of the light particles and hence it is stable for the even m (and even p). In the case of $m = \text{odd}$ ($p = \text{odd}$), the $\mathcal{O}$ and $\mathcal{O}'$ must contain odd total number of matter multiplets, and thus the X particle with even R-parity is stable as shown in the previous $\mathbf{Z}_6$ and $\mathbf{Z}_{18}$ models.

We now see that the relaxed anomaly-free condition (8) still plays an important role to prove the stability of the superheavy X particle. (A crucial point is that the $\mathbf{Z}_N$ charges p is even, if and only if the m is even.)

If the m (and p) is odd and the X has the R-parity opposite to that in the stable case, it can decay into the SUSY standard-model particles. Even in this case, however, the lifetime of the X particle can be longer than the age of the

universe depending on the choice of N, p and m . In fact, for $N = 10$ (i.e. $\mathbf{Z}_{10}$)⁵ we have found no nonrenormalizable operator $\mathcal{O}$ and $\mathcal{O}'$ making the X particle to decay up to the eighth order. In the ninth order we find a possible superpotential

$$W = \left(\frac{1}{M_*}\right)^7 \bar{d}\bar{d}\bar{d}lllH_u lH_u X, \quad (10)$$

in the $\mathbf{Z}_{10}$ with $(p, m) = (1, 3)$ and $(3, 9)$. The lifetime of the X particle is of the order of

$$\tau_X \sim \left(\frac{M_*}{M_X}\right)^{14} \frac{1}{M_X} \simeq 10^{11} - 10^{26} \text{ years}, \quad (11)$$

for the cut-off scale $M_* \simeq 10^{18}$ GeV and $M_X \simeq 10^{13} - 10^{14}$ GeV. This is enough long for X to be a perfect superheavy dark matter. The lowest dimensional operator and the lifetime of X in each model, $(N, p$ and $m)$, are also given for $N \leq 18$ in Table 4.

More interestingly, if the X particles are unstable, their decays might explain the ultra-high-energy cosmic ray events [13] observed recently beyond the Greisen-Zatsepin-Kuzmin (GZK) cut-off energy [14]. If the X particles are the dark matter in the galaxy, the observed air showers may be mostly due to gamma rays and nucleons produced by the X decays in the halo of our galaxy and they can easily exceed the GZK cut-off energy if $M_X \gtrsim 10^{13}$ GeV [6, 8]. If the X particles form a dominant part of the dark matter, we may fit the observed fluxes with the lifetime $10^{15} \text{ years} \lesssim \tau_X \lesssim 10^{22} \text{ years}$ [9, 4, 15]. We should stress that the X particle with odd R-parity in the $\mathbf{Z}_{10}$ models meets nicely this condition (see eq. (11)). It may be very encouraging that these $\mathbf{Z}_{10}$ models can be embedded into the grand unified $\text{SU}(5)_{GUT}$.⁶

So far we have assumed X to be a singlet of the standard-model gauge groups. However, our conclusion can be also applied to the case of X being $\mathbf{8}$ of $\text{SU}(3)_C$ or $\mathbf{3}$ of $\text{SU}(2)_L$. It may be interesting that these particles with masses of the order of $10^{13} - 10^{14}$ GeV may resolve the problem of apparent discrepancy between the gauge-coupling unification and the string scales [16].

⁵ For the $\mathbf{Z}_{10}$ model to be anomaly free, we must take the Kac-Moody level $k = 5$ in eq. (8) for the standard-model gauge groups $\text{SU}(3)_C$ and $\text{SU}(2)_L$.

⁶ By using a $\text{U}(1)_Y$ gauge rotation one can make the $\mathbf{Z}_N$ charges for all light particles to be consistent with $\text{SU}(5)_{GUT}$.

N	(p, m)	k	the lowest dimensional operator	R-parity	lifetime
4	(1, 1)	16	$lH_u lH_u \quad (\in \mathcal{O})$	+	$10^{-22} - 10^{-17}$ sec
8	(1, 5)	32	$\bar{d}\bar{d}\bar{d}ll lH_u \quad (\in \mathcal{O})$	+	$10^2 - 10^{13}$ sec
8	(3, 7)	32	$\bar{d}\bar{d}\bar{d}ll lH_u \quad (\in \mathcal{O})$	+	$10^2 - 10^{13}$ sec
10	(1, 3)	5	$\bar{d}\bar{d}\bar{d}ll lH_u lH_u \quad (\in \mathcal{O})$	-	$10^{11} - 10^{26}$ years
10	(3, 9)	5	$\bar{d}\bar{d}\bar{d}ll lH_u lH_u \quad (\in \mathcal{O})$	-	$10^{11} - 10^{26}$ years
12	(1, 1)	16	$\bar{u}\bar{u}\bar{d}\bar{d}\bar{d}\bar{d} \quad (\in \mathcal{O})$	+	$10^{-6} - 10^3$ sec
12	(5, 1)	16	$\bar{u}\bar{u}\bar{d}\bar{d}\bar{d}\bar{d} \quad (\in \mathcal{O})$	+	$10^{-6} - 10^3$ sec
14	(1, 9)	7	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u \quad (\in \mathcal{O})$	-	$10^{35} - 10^{56}$ years
14	(3, 13)	7	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u \quad (\in \mathcal{O})$	-	$10^{35} - 10^{56}$ years
14	(5, 3)	7	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u \quad (\in \mathcal{O})$	-	$10^{35} - 10^{56}$ years
16	(1, 5)	64	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u lH_u \quad (\in \mathcal{O})$	+	$10^{51} - 10^{76}$ years
16	(3, 15)	64	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u lH_u \quad (\in \mathcal{O})$	+	$10^{51} - 10^{76}$ years
16	(5, 9)	64	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u lH_u \quad (\in \mathcal{O})$	+	$10^{51} - 10^{76}$ years
16	(7, 3)	64	$\bar{d}\bar{d}\bar{d}ll \bar{d}\bar{d}\bar{d}ll lH_u lH_u \quad (\in \mathcal{O})$	+	$10^{51} - 10^{76}$ years

Table 4: The lowest dimensional operators and the lifetimes of the superheavy X particle in the anomaly-free $\mathbf{Z}_N$ models with $N \leq 18$. Here, we have assumed the mass of X , $M_X \simeq 10^{13} - 10^{14}$ GeV. k denotes the Kac-Moody levels of the $SU(3)_C$ and $SU(2)_L$ gauge groups. For $N = 6$ and 18, see Table 2.

References

- [1] K. Griest and M. Kamionkowski, *Phys. Rev. Lett.* **64** (1990) 615.
- [2] J. Ellis, J.L. Lopez, and D.V. Nanopoulos, *Phys. Lett.* **B247** (1990) 257.
- [3] D.J.H. Chung, E.W. Kolb, and A. Riotto, hep-ph/9802238.
- [4] V. Kuzmin and I. Tkachev, hep-ph/9802304.
- [5] S. Coleman, *Nucl. Phys.* **B307** (1988) 867.
- [6] V. Berezhinsky, M. Kachelriess, and A. Vilenkin, *Phys. Rev. Lett.* **79** (1997) 4302.
- [7] L.M. Krauss and F. Wilczek, *Phys. Rev. Lett.* **62** (1989) 1221.
- [8] V.A. Kuzmin and V.A. Rubakov, astro-ph/9709187.
- [9] K. Benakli, J. Ellis, and D.V. Nanopoulos, hep-ph/9803333.
- [10] L.E. Ibanez and G.G. Ross, *Nucl. Phys.* **B368** (1992) 3.
- [11] M.B. Green and J.H. Schwarz, *Phys. Lett.* **B149** (1984) 117.
- [12] L.E. Ibanez, *Nucl. Phys.* **B398** (1993) 301.
- [13] N. Hayashida et.al., *Phys. Rev. Lett.* **73** (1994) 3491;
D.J. Bird et.al., *sl Astrophys. J.* **424** (1994) 491;
T.A. Egorov et.al., in: *Proc. Tokyo Workshop on Techniques for the Study of Extremely High Energy Cosmic Rays*, ed. M. Nagano (ICRR, U. of Tokyo, 1993).
- [14] K. Greisen, *Phys. Rev. Lett.* **16** (1966) 748;
G.T. Zatsepin and V.A. Kuzmin, *Pisma Zh. Eksp. Theor. Fiz.* **4** (1966) 114.
- [15] M. Birkel and S. Sarkar, hep-ph/9804285.
- [16] C. Bachas, C. Fabre, and T. Yanagida, *Phys. Lett.* **B370** (1996) 49;
T. Han, T. Yanagida, and R. Zhang, hep-ph/9804228.